



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Polyporeae.....	M. A. Levine
Exobasidii.....	H. M. Richards
Rusts and Smuts.....	E. W. Olive
Discomycetes.....	B. O. Dodge
Lichens.....	W. C. Barbour
Pyrenomycetes, Sphaeriaceae and Dothideaceae.....	H. M. Richards
Hypocreaceae, Perisporieae, Plectascineae, Tuberineae,	
	F. J. Seaver
Fungi forming Sclerotia.....	A. B. Stout
Imperfecti.....	H. M. Richards, F. J. Seaver, Mel T. Cooke
Oomycetes.....	C. A. King
Zygomycetes.....	A. F. Blakeslee
Chytridiaceae.....	
Myxomycetes.....	Miss M. E. Latham
Yeasts and Bacteria.....	Miss J. Broadhurst
Insect Galls.....	Mel T. Cooke

TWO NEW PLANTS FROM THE TERTIARY ROCKS OF THE WEST

BY T. D. A. COCKERELL

***Smilax labidurommæ* sp. nov.**

Leaf 53 mm. long and 35 wide; deltoid, with truncate base, the lateral margins nearly straight, but under a lens showing shallow crenulation; five principal longitudinal veins.

Miocene shales, Florissant, Colorado, Station 14 (*Wilmatte P. Cockerell*). On the same slab, three mm. from the leaf, is an earwig, *Labiduromma bormansi* Scudder. The genus *Smilax* and the family Smilacæ are new to the Florissant list, but various species of *Smilax* have been found in other American formations. ***Smilax carbonensis*** n. n. (*S. grandifolia* Lesq., Tertiary Flora, Pl. IX, f. 5, from Carbon, Wyoming) is a larger leaf, with cordate instead of truncate base, and convex lateral margins. It is probably quite distinct from *Smilacites grandifolia* Unger,* which as originally figured by that author, has the basal sinus very deep (over 30 mm.); and in any event *S. grandifolia* Buck-

* *Chloris Protogaea*, pl. XL, f. 3.

ley* antedates Unger's name by about four years. In the determination of *Smilax* leaves there is indeed a large element of uncertainty, owing to the variation in outline, as Laurent† has beautifully illustrated in the case of *S. aspera*. This should prevent us from multiplying specific names based on different




FIG. 1. *Smilax labidurommæ* Cockerell.

looking leaves of the same region and period, but on the other hand, it should not lead us to consider identical plants of quite different parts of the world and different geological horizons. Something must be allowed for the inherent probabilities in each case. Knowlton‡ has described *Smilax lamarensis* from the supposed Miocene of the Yellowstone; it resembles *S. carbonensis* rather than the Florissant species, having the cordate base and rounded sides. Heer,§ from beds supposed to be Miocene at Asakak, Greenland, describes a *Smilax lingulata*; it is a narrow leaf quite unlike the Florissant plant, resembling, in fact, the living *S. laurifolia* L.

A much more ancient plant assigned to this genus is **Smilax**

* Am. Journ. Sci. 45: 171. 1843.

† Ann. Mus. Marseille, 12: pl. 1. (1908.)

‡ Geol. Yellowstone Nat. Park, pl. CXXI, f. 3, 4.

§ Kongl. Sv. Vet. Akad. Handl., 13, no. 2: 15, pl. 1, f. 12 (1874).

Kansana n. n. (*S. undulata* Lesq.,* not of Pohl†), from the Dakota group in Kansas.

***Tithymalus phenacodorum* sp. nov.**

Seed. Length 4.75, breadth 4.25 mm.; short-pyriform, with four sides slightly flattened; surface coarsely irregularly wrinkled.

Five miles southeast of mouth of Pat O'Hara Creek, Clark's Fork Basin, Wyoming; above red-banded beds, in strata supposed to be older than the Wasatch, though formerly classed



FIG. 2. Seeds of *Tithymalus*: A. *T. phenacodorum*, side view. B. *T. phenacodorum*, from above. C. *T. Willistoni*, side view.

with that group. Type in American Museum of Natural History; collected by Mr. W. Stein, along with numerous remarkable land shells, of the genera *Protoboysia*, *Boysia*, *Vitrea*, *Thysanophora*, *Pyramidula*, *Gastrodonta* and *Oreohelix*.

Compared with the seeds of *T. Willistoni* Cockerell,‡ from the Loup Fork Beds of Long I., Kansas, the new species is distinctly longer, while the depressions between the rugæ are more irregular and less definitely in longitudinal series. A quite similar but still longer seed is that of the living Mexican ***Tithymalus campestris*** n. n. (*Euphorbia campestris* Cham. & Schl., 1830), the seed of which is well figured by Millspaugh, Botanical Gazette, XXV (1898): 25. In *T. campestris*, however, the rugæ are more labyrinthiform than in *T. phenacodorum*.

Extremely hard and dense seeds, such as those of *Tithymalus*, are readily fossilized where ordinary vegetation decays and disappears. It is probable that careful search will reveal them in other Tertiary strata. I add a figure of *T. Willistoni*, as it has not been figured.

* Fl. Dakota Group: 39 (1892).

† Pohl, A. DC. in D. C., Mon. Phan. 1: 135.

‡ Torrey 9: 119. 1909.